

Topological insulators with SU(2) Landau levels

Yi Li,¹ Shou-Cheng Zhang,² and Congjun Wu^{1,3}

¹*Department of Physics, University of California, San Diego, La Jolla, California 92093, USA*

²*Department of Physics, Stanford University, Stanford, California 94305, USA*

³*Key Laboratory of Artificial Micro- and Nano-structures of Ministry of Education, and School of Physics and Technology, Wuhan University, Wuhan 430072, China*

We construct continuum models of 3d and 4d topological insulators by coupling spin- $\frac{1}{2}$ fermions to an SU(2) background gauge field, which is equivalent to a spatially dependent spin-orbit coupling. Higher dimensional generalizations of flat Landau levels are obtained in the Landau-like gauge. The 2d helical Dirac modes with opposite helicities and 3d Weyl modes with opposite chiralities are spatially separated along the third and fourth dimensions, respectively. Stable 2d helical Fermi surfaces and 3d chiral Fermi surfaces appear on open boundaries, respectively. The charge pumping in 4d Landau level systems shows quantized 4d quantum Hall effect.

PACS numbers: 73.43.Cd, 71.70.Ej, 73.21.-b

Time-reversal (TR) invariant topological insulators (TI) have become a major research focus in condensed matter physics[1–3]. Different from the 2d quantum Hall (QH) and quantum anomalous Hall systems which are topologically characterized by the first Chern number [4–6], time-reversal invariant TIs are characterized by the second Chern number in 4d [7, 8] and the Z_2 index in 2d and 3d[8–14]. Various 2d and 3d TIs are predicted theoretically and identified experimentally exhibiting stable gapless 1D helical edge and 2d surface modes against TR invariant perturbations [11, 15–19]. Topological states have also been extended to systems with particle-hole symmetry and superconductors [20–22].

Most current studies of 2d and 3d TIs focuses on Bloch-wave bands in lattice systems. The non-trivial band topology arises from spin-orbit (SO) coupling induced band inversions[11]. However, Landau levels (LL) are essential in the study of QH effects because their elegant analytical properties enable construction of fractional states. Generalizing LLs to high dimensions gives rise to TIs with explicit wavefunctions in the continuum, which would facilitate the study of the exotic fractional TIs. Efforts along this line were pioneered by one of the authors and Hu [7]. They constructed LLs on the compact S^4 sphere by coupling fermions with the SU(2) monopole gauge potential, and various further developments appeared [23–27]. 2d TIs based on TR invariant LLs have also been investigated [10]. Two of the authors generalized LLs of Schrödinger fermions to high-dimensional flat space [28] by combining the isotropic harmonic potential and SO coupling. LLs have also been generalized to high dimensional Dirac fermions and parity-breaking systems [29, 30].

In all the above works, angular momentum is explicitly conserved, thus they can be considered as LLs in the symmetric-like gauge. In 2d, LL wavefunctions in the Landau gauge are particularly intuitive: they are 1d chiral plane-wave modes spatially separated along the transverse direction. The QH effect is just the 1d chiral

anomaly in which the chiral current generated by the electric field becomes the transverse charge current. In this article, we develop high dimensional LLs with flat spectra as spatially separated helical Dirac or chiral Weyl fermion modes, i.e., the SU(2) Landau-like gauge. They are 3d and 4d TIs defined in the continuum possessing stable gapless boundary modes. For the 4d case, they exhibit the 4d quantum Hall effect[7, 8], which is a quantized non-linear electromagnetic response related to the spatially separated (3+1)d chiral anomaly. Our methods can be easily generalized to arbitrary dimensions and also to Dirac fermions.

The 3d case We begin with the 3d TR invariant LL Hamiltonian for a spin- $\frac{1}{2}$ fermion as

$$H_{LL}^{3d,\nu=\pm} = \frac{\vec{p}^2}{2m} + \frac{1}{2}m\omega_{so}^2 z^2 - \nu\omega_{so}z(p_x\sigma_y - p_y\sigma_x)(1)$$

which couples the 1D harmonic potential in the z direction and the 2d Rashba SO coupling through a z -dependent SO coupling strength. $H_{LL}^{3d,\nu}$ possess translation symmetry in the xy -plane, TR and parity symmetries. $H_{LL}^{3d,+}$ and $H_{LL}^{3d,-}$ share the same physics up to a $z \rightarrow -z$ transformation. For simplicity, unless explicitly mentioned, we only consider $H_{LL}^{3d,+}$ and suppress the index ν . Eq. 1 can be reformulated in the form of an SU(2) background gauge potential as $H_{LL}^{3d} = \frac{1}{2m}(\vec{p} - \frac{e}{c}\vec{A})^2 - \frac{1}{2}m\omega_{so}^2 z^2$, where $\omega_{so} = |eG|/mc$, and $\vec{A}$ takes the Landau-like gauge as $A_x = G\sigma_y z$, $A_y = -G\sigma_x z$ and $A_z = 0$. In Ref. 28, a symmetric-like gauge with $\vec{A}' = G\vec{\sigma} \times \vec{r}$ is used, which explicitly preserves the 3d rotational symmetry. However, the SU(2) vector potentials $\vec{A}'$ and $\vec{A}$ are *not* gauge equivalent. As shown below, the physical quantities of Eq. 1, such as density of states (DOS), are not 3d rotationally symmetric. Nevertheless, these two Hamiltonians belong to the same topological class. A related Hamiltonian is also employed for studying electromagnetic properties in superconductors with cylindrical geometry [31].

Eq. (1) can be decomposed into a set of 1D harmonic

oscillators along the z -axis exhibiting flat spectra, a key feature of LLs. We define a characteristic SO length scale $l_{so} = \sqrt{\frac{\hbar}{m\omega_{so}}}$. Each of the reduced 1D harmonic oscillator Hamiltonian is associated with a 2d helical plane-wave state as $H^z(\vec{k}_{2d}) = \frac{p_z^2}{2m} + \frac{1}{2}m\omega_{so}^2[z - l_{so}^2 k_{2d} \hat{\Sigma}_{2d}(\hat{k}_{2d})]^2$, where $k_{2d} = (k_x^2 + k_y^2)^{\frac{1}{2}}$ and $\hat{k}_{2d} = (k_x, k_y)$; the helicity operator is defined as $\hat{\Sigma}_{2d}(\hat{k}_{2d}) = \hat{k}_x \sigma_y - \hat{k}_y \sigma_x$. The n -th LL eigenstates are solved as

$$\Psi_{n, \vec{k}_{2d}, \Sigma}(\vec{r}) = e^{i\vec{k}_{2d} \cdot \vec{r}_{2d}} \phi_n[z - z_0(k_{2d}, \Sigma)] \otimes \chi_{\Sigma}(\hat{k}_{2d}), \quad (2)$$

where $\vec{r}_{2d} = (x, y)$; $\chi_{\Sigma}(\hat{k}_{2d})$ are eigenstates of the helicity satisfying $\hat{\Sigma} \chi_{\Sigma}(\hat{k}_{2d}) = \Sigma \chi_{\Sigma}(\hat{k}_{2d})$ with helicity eigenvalues $\Sigma = \pm 1$; $\phi_n[z - z_0]$ are the eigenstates of the n -th harmonic levels with the central positions located at z_0 , and $z_0(k_{2d}, \Sigma) = \Sigma l_{so}^2 k_{2d}$. The energy spectra of the n -th LL is $E_n = (n + \frac{1}{2})\hbar\omega_{so}$, independent of $\vec{k}_{2d}$ and Σ .

The classical equations of motion for H_{LL}^{3d} are derived as

$$\begin{aligned} \dot{\vec{r}} &= \frac{\vec{p} - \frac{e}{c}\vec{A}}{m}, \quad \dot{p}_{2d} = 0, \quad \dot{p}_z = 2\omega(\vec{p} \times \vec{S})_z - m\omega^2 z, \\ \dot{\vec{S}}_{2d} &= -\frac{2\omega}{\hbar} z S_z \vec{p}_{2d}, \quad \dot{S}_z = \frac{2\omega}{\hbar} z \vec{S}_{2d} \cdot \vec{p}_{2d}, \end{aligned} \quad (3)$$

where $\vec{S}_{2d} = (\frac{1}{2}\sigma_x, \frac{1}{2}\sigma_y)$ and $S_z = \frac{1}{2}\sigma_z$. If we choose the initial condition of spin $\vec{S}$ such that $S_{z,0} = 0$ and $\vec{S}_{2d,0} \perp \vec{p}_{2d,0}$, then $\vec{S}$ is conserved and lies in the xy -plane. The motion is reduced to a coplanar cyclotron one in the vertical plane perpendicular to $\vec{S}$. In other words, the orbital angular momentum and spin are locked, a feature from SO coupling.




FIG. 1: (Color online) 3d and 4d LLs for H_{LL}^{3d} and H_{LL}^{4d} as spatially separated 2d helical SO plane-wave modes localized along the z -axis (A), and 3d Weyl modes localized along the u -axis (B), respectively. Their central locations are $z_0(\vec{k}_{2d}, \Sigma) = \Sigma l_{so}^2 k_{2d}$ and $u_0(\vec{k}_{3d}, \Sigma) = \Sigma l_{so}^2 k_{3d}$, respectively. Note that 2d Dirac modes with opposite helicities and the 3d ones with opposite chiralities are located at opposite sides of $z = 0$ and $u = 0$ planes, respectively.

In the 2d LL case, the spatial coordinates x and y are non-commutative if projected to a given LL, say, the lowest LL (LLL). The LLL wavefunctions are chiral 1d plane-waves along the x -direction whose central

y -position is $y_0 = l_B^2 k_x$, thus 1d chiral modes with opposite chiralities are spatially separated. Consequently, the xy -plane can be viewed as the 2D phase space of a 1d system, in which y plays the role of k_x . The momentum cut-off of the bulk states is determined by the system size along the y -direction as $|k_x| < \frac{L_y}{2l_B^2}$. Similarly, the 3d LL wavefunctions in Eq. 2 are spatially separated 2d helical plane-waves along the z -axis. As shown in Fig. 1 (A), for states with opposite helicity eigenvalues, their central positions are shifted in opposite directions. After the projection into the LLL, z is equivalent to $l_{so}^2(k_x \sigma_y - k_y \sigma_x)$, and thus

$$[x, z]_{LLL} = il_{so}^2 \sigma_y, [y, z]_{LLL} = -il_{so}^2 \sigma_x, [x, y]_{LLL} = 0. \quad (4)$$

Interestingly, the 3d LL states can be viewed as states in the 4D phase space of a 2d system (with x and y coordinates), in which $|z|$ plays the role of the magnitude of the 2d momentum. In fact, the momentum cut-off of the bulk states is exactly determined by the spatial size L_z along the z direction as

$$k_{2d} < k_{2d}^B \equiv \frac{L_z}{2l_{so}^2}, \quad (5)$$

in which $-\frac{L_z}{2} < z < \frac{L_z}{2}$. Applying the open boundary condition along the z direction, and periodic boundary conditions in the x and y direction with spatial sizes L_x and L_y respectively, we can easily count the total number of states to be $\mathcal{N} = \frac{L_x L_y L_z}{8\pi l_{so}^4}$. The L_z^2 dependence of $\mathcal{N}$ may seem puzzling for a 3d system, but expressing L_z in terms of Eq. (5), we find $\mathcal{N} = \frac{1}{2\pi} L_x L_y (k_{2d}^B)^2$, which is the conventional state counting of a 2d system expressed in terms of the 4D phase space volume.

The topological property of this 3d LL system manifests through its helical surface spectra. Let us consider the upper boundary located at $z = z_B$ for H_{LL}^{3d} . For simplicity, we only consider the LLL as an example. If $z_B > 0$, the positive helicity states $\Psi_{0, \vec{k}_{2d}, \Sigma=1}$ with $k_{2d} > k_{2d}^B = z_B l_{so}^{-2}$ are confined at the boundary. The 1d harmonic potential associated with $\vec{k}_{2d}$ and $\Sigma = 1$ is truncated at $z = z_B$, and thus the surface spectra acquire dispersion. If we neglect the zero-point energy, the surface mode dispersion is approximated by $\frac{1}{2}m\omega_{so}^2(k - k_{2d}^B)^2 l_{so}^4$ with $k > k_{2d}^B$. If the chemical potential μ lies above the LLL, it cuts the spectra at surface states with a Fermi wavevector $k_f > k_{2d}^B$. The Fermi velocity is $v_f \approx m(k_f - k_{2d}^B)\omega_{so}^2 l_{so}^4$. The surface Hamiltonian is approximated as $H_{sf} \approx v_f(\vec{p} \times \vec{\sigma}) \cdot \hat{z} - \mu$ with an electron-like Fermi surface with $\Sigma = 1$. In another case, if the upper boundary is located at $z_B < 0$, then the negative helicity states $\Psi_{0, \vec{k}_{2d}, \Sigma=-1}$ with $k_{2d} < k_{2d}^B$, and all the positive helicity states are pushed to the boundary as surface modes. Depending on the value of μ , we can have a hole-like Fermi surface with $\Sigma = -1$, a Dirac Fermi point, or, an electron-like Fermi surface with $\Sigma = 1$. Similarly,

any other LL gives rise to a branch of gapless helical surface modes, and each filled bulk LL contributes one helical Fermi surface. For the lower boundary, the analysis is parallel to the above. Each filled LL gives rise to an electron-like helical Fermi surface with $\Sigma = -1$, or hole-like with $\Sigma = 1$.

The 4d case The above procedure can be straightforwardly generalized to any higher dimension. For example, the 4d LL Hamiltonian is denoted as

$$H_{LL}^{4d,\nu=\pm} = \frac{p_u^2}{2m} + \frac{1}{2}m\omega^2 u^2 + \frac{\vec{p}_{3d}^2}{2m} - \nu\omega u \vec{p}_{3d} \cdot \vec{\sigma}, \quad (6)$$

where u and p_u refer to the coordinate and momentum of the 4th dimension, respectively, and $\vec{p}_{3d}$ is the 3-momentum in the xyz -space. Without loss of the generality, let us focus on the case of $\nu = +$ and Eq. 6 can be represented as $H_{LL}^{4d} = \frac{1}{2m} \sum_{i=1}^4 (p_i - \frac{e}{c} A_i)^2 - m\omega^2 u^2$, where the SU(2) vector potential takes the Landau-like gauge with $A_i = G\sigma_i u$ for $i = x, y, z$ and $A_u = 0$. Eq. 6 preserves the translational and rotational symmetries in the xyz -space and TR symmetry. Similar to the 3d case, the 4d LL spectra and wavefunctions are solved by reducing Eq. 6 into a set of 1d harmonic oscillators along the u -axis as $H^u(\vec{k}_{3d}) = \frac{p_u^2}{2m} + \frac{1}{2}m\omega^2(u - l_{so}^2 k_{3d} \hat{\Sigma}_{3d})^2$ where $k_{3d} = (k_x^2 + k_y^2 + k_z^2)^{\frac{1}{2}}$ and $\hat{\Sigma}_{3d} = \hat{k}_{3d} \cdot \vec{\sigma}$. The LL wavefunctions are

$$\Psi_{n,\vec{k}_{3d},\Sigma}(\vec{r}, u) = e^{i\vec{k}_{3d} \cdot \vec{r}} \phi_n[u - u_0(k_{3d}, \Sigma)] \otimes \chi_{\Sigma}(\vec{k}_{3d}), \quad (7)$$

where, the central position $u_0(k_{3d}, \Sigma) = \Sigma l_{so}^2 k_{3d}$; χ_{Σ} are eigenstates of 3d helicity $\hat{\Sigma}_{3d}$ with eigenvalues $\Sigma = \pm 1$. Inside each LL, the spectra are flat with respect to $\vec{k}_{3d}$ and Σ . This realizes the spatial separation of the 3d Weyl fermion modes as shown in Fig. 1 (B). After the LLL projection for H_{LL}^{4d} , the non-commutative relations among coordinates are $[r_i, u] = il_{so}^2 \sigma_i$ and $[r_i, r_j] = 0$ for $i = x, y, z$. Similar to the discussions of the 3d case, the 4d LLs can be viewed as states of the 6D phase space of a 3d system: increasing the width along the u -direction corresponds to increasing the bulk momentum cut-off $k_{3d} < k_{3d}^B \equiv L_u/(2l_{so}^2)$. If the LLL is fully-filled, the total number of states is given by $\mathcal{N} = \frac{L_x L_y L_z L_u^3}{24\pi^2 l_{so}^6}$. Re-expressing $L_u = 2k_{3d}^B l_{so}^2$ we find $\mathcal{N} = \frac{1}{3\pi^2} L_x L_y L_z (k_{3d}^B)^3$, which is the conventional state counting of a 3d system expressed in terms of the 6D phase space volume. With an open boundary imposed along the u direction, 3d helical Weyl fermion modes appear on the boundary.

Now let us consider the generalized 4d QHE[7, 8] as the nonlinear electromagnetic response of $(4+1)$ d LL system to the external electric and magnetic (EM) fields, with $\vec{E} \parallel \vec{B}$ in the xyz -space. Without loss of generality, we choose the EM fields as $\vec{E} = E\hat{z}$ and $\vec{B} = B\hat{z}$, which are minimally coupled to the spin-1/2 fermion,

$$H_{LL}^{4d}(E, B) = -\frac{\hbar^2}{2m} \nabla_u^2 + \frac{1}{2}m\omega^2 \left(u + il_{so}^2 \vec{D} \cdot \vec{\sigma} \right)^2, \quad (8)$$




FIG. 2: The central positions $u_0(m, k_z, \nu)$ of the 4d LLs in the presence of the magnetic field $\vec{B} = B\hat{z}$. The branch of $m = 0$ runs across the entire u -axis, which gives rise to quantized charge transport along u -axis in the presence of $\vec{E} \parallel \vec{B}$ as indicated in Eq. 10.

where $\vec{D} = \vec{\nabla} - i\frac{e}{\hbar c} \vec{A}_{em}$. Here $\vec{A}_{em}$ is the $U(1)$ magnetic vector potential in the Landau gauge with $A_{em,x} = 0$, $A_{em,y} = Bx$ and $A_{em,z} = -cEt$. We define $l_B = \sqrt{\frac{\hbar c}{eB}}$, where $eB > 0$ is assumed.

The $\vec{B}$ -field further reorganizes the chiral plane-wave states inside the n -th 4d LL into a series of 2d magnetic LLs in the xy -plane. For the moment, let us set $E_z = 0$. The eigenvalues $E_n = (n + \frac{1}{2})\hbar\omega_{so}$ remain the same as before without splitting, while the eigen-wavefunctions are changed. We introduce a magnetic LL index m in the xy -plane. For the case of $m = 0$, the eigen-wavefunctions are spin polarized as

$$\begin{aligned} \Psi_{n,m=0}(k_y, k_z) &= e^{ik_y y + ik_z z} \phi_n[u - u_0(k_z, m=0)] \\ &\otimes \chi_{m=0}[x - x_0(k_y)], \end{aligned} \quad (9)$$

where $x_0 = l_B^2 k_y$ and $\chi_0 = [\phi_0(x - x_0), 0]^T$ is the zero mode channel of the operator $-i\vec{D} \cdot \vec{\sigma}$ with the eigenvalue of $\lambda_0 = k_z$. The central positions of the u -direction harmonic oscillators are $u_0(k_z, m=0) = l_{so}^2 k_z$. For $m \geq 1$, the eigen-modes of $-i\vec{D} \cdot \vec{\sigma}$ come in pairs as $\chi_{m,\pm}[x - x_0(k_y)] = [\alpha_{m,\pm}\phi_m(x - x_0), \beta_m\phi_{m-1}(x - x_0)]^T$, where, coefficients $\alpha_{m,\pm} = l_B k_z \pm \sqrt{l_B^2 k_z^2 + 2m}$, $\beta_m = -i\sqrt{2m}$, and the eigenvalues are $\lambda_{m,\pm} = \pm\sqrt{k_z^2 + 2ml_B^{-2}}$. The corresponding eigen-wavefunctions are $\Psi_{n,m,\pm}(k_y, k_z) = e^{ik_y y + ik_z z} \phi_n[u - u_0(k_z, m, \pm)] \chi_{m,\pm}[x - x_0(k_y)]$, where the central positions $u_0(k_z, m, \pm) = \pm l_{so}^2 \sqrt{k_z^2 + 2ml_B^{-2}}$.

For the solutions of Eq. 9 with the same 4d LL index n , the 2d magnetic LL with index $m = 0$ is singled out. The central positions of states in this branch is linear with k_z , and thus run across the entire u -axis, while those of other branches with $m \geq 1$ only lie in one half of the space as shown in Fig. 2. After turning on E_z , k_z is accelerated with time as $k_z(t) = k_z(0) + eE_z t/\hbar$, and thus the central position $u_0(m=0, k_z)$ moves along the u -axis. Only the $m = 0$ branch of the magnetic LL states contribute to the charge pumping which results in an electric current along the u -direction. Within the time interval Δt , the number of states with each filled 4d LL passing a cross

section perpendicular to the u -axis is $N = \frac{L_x L_y}{2\pi l_B^2} \frac{eE\Delta t L_z}{2\pi}$, which results in the electric current density $\frac{e}{L_x L_y L_z} \frac{dN}{dt}$. If the number of fully filled 4d LLs is n_{occ} , the total current density along the u -axis is

$$j_u = n_{occ} \alpha \frac{e}{4\pi^2 \hbar} \vec{E} \cdot \vec{B}, \quad (10)$$

where $\alpha = e^2/(\hbar c)$ is the fine-structure constant. This quantized non-linear electromagnetic response is in agreement with results from the effective theory [8] as the 4d generalization of the QH effect. If we impose open boundary conditions perpendicular to the u -direction, the above charge pump process corresponds to the chiral anomalies of Weyl fermions with opposite chiralities on the two 3d boundaries, respectively. Since they are spatially separated, the chiral current corresponds to the electric current along the u -direction.

Although the DOS of our 4d LL systems is different from the 4d TIs in the lattice system with translation symmetry [8], their electromagnetic responses obey the same Eq. 10. It is because that only the spin-polarized $m = 0$ branch of LLs, $\Psi_{n,m=0}(k_y, k_z)$, is responsible for the charge pumping. For this branch, B_z -field quantizes the motion in the xy -plane so that the x and u coordinates play the role of k_y and k_z , respectively. Consequently, the system can be viewed as the 4D phase space of coordinates y and z , and scales uniformly as $L_x L_y L_z L_u$ with the conventional thermodynamic limit of a 4d system.

High dimensional LLs of Dirac electrons The above procedure can be generalized to Dirac fermions as square root problems to Eq. (1) and Eq. (6). For the 3d case, we have

$$H_{LL,Dirac}^{3d} = \frac{l_{so}\omega_{so}}{\sqrt{2}} \begin{bmatrix} 0 & \vec{\sigma} \cdot \vec{p} + i \frac{z\hbar}{l_{so}^2} \sigma_z \\ \vec{\sigma} \cdot \vec{p} - i \frac{z\hbar}{l_{so}^2} \sigma_z & 0 \end{bmatrix}, \quad (11)$$

whose square exhibits a diagonal block form with a supersymmetric structure as $(H_{LL,Dirac}^{3d})^2/(\hbar\omega_{so}) = \text{diag}(H_{LL}^{3d,+} - \frac{\hbar\omega_{so}}{2}, H_{LL}^{3d,-} + \frac{\hbar\omega_{so}}{2})$. The eigenvalues of Eq. (11) are $E_{\pm n} = \pm \sqrt{n} \hbar \omega_{so}$. The eigen-wavefunctions are constructed based on the eigenstates of $H_{LL}^{3d,\nu=\pm}$ as $\Psi_{\pm n,Dirac}^{3d}(\vec{k}_{2d}, \Sigma) = \frac{1}{\sqrt{2}} [\Psi_{n,\vec{k}_{2d},\Sigma}^+ \pm \Psi_{n-1,\vec{k}_{2d},-\Sigma}^-]^T$, where $\Psi_{n,\vec{k}_{2d},\Sigma}^- = e^{i\vec{k}_{2d} \cdot \vec{r}_{2d}} \phi_n[z + z_0(k_{2d}, \Sigma)] \otimes \chi_\Sigma(k_{2d})$ is the eigenstate of $H_{LL}^{3d,-}$. The 0th LL states are Jackiw-Rebbi half-fermion modes with only the upper two components nonzero [32, 33]. The 4d Hamiltonian can be constructed as

$$H_{LL,Dirac}^{4d} = \frac{l_{so}\omega_{so}}{\sqrt{2}} \begin{bmatrix} 0 & \vec{\sigma} \cdot \vec{p}_{3d} - i \frac{a_u}{l_{so}} \\ \vec{\sigma} \cdot \vec{p}_{3d} + i \frac{a_u}{l_{so}} & 0 \end{bmatrix}, \quad (12)$$

where $a_u = \frac{1}{\sqrt{2}l_{so}}(u + i \frac{l_{so}^2}{\hbar} p_u)$ is the phonon annihilation operator in the u -direction. The eigen-values

are still $E_{\pm n} = \pm \sqrt{n} \hbar \omega_{so}$, and the eigenstates are $\Psi_{\pm n,Dirac}^{4d}(\vec{k}_{3d}, \Sigma) = \frac{1}{\sqrt{2}} [\Psi_{n,\vec{k}_{2d},\Sigma}^+ \pm \Psi_{n-1,\vec{k}_{2d},\Sigma}^-]^T$.

Laughlin-type wavefunctions The construction of the Laughlin-type wavefunctions for interacting fermions is difficult for these SO coupled high dimensional LL systems. Nevertheless, for the 4d case in the magnetic field, the LLL states with both $n = 0$ and $m = 0$ are spin-polarized, and their total DOS is finite as $\rho_{n=m=0}^{4d} = \frac{1}{4\pi^2 l_{so}^2 l_B^2}$. Even though they are degenerate with other LLL states with $(n = 0, m \neq 0)$, they are favored by repulsive interactions if they are partially filled. The Laughlin wavefunction in the Landau gauge for the 2d LLs has been constructed in Ref. [34]. We generalize it to the 4d case, and define $w = e^{i \frac{x+iy}{L_x}}$ and $v = e^{i \frac{z+iu}{L_z}}$, then LL states are represented as $w^{h_x} v^{h_z}$ up to a Gaussian factor $e^{-\frac{y^2}{2l_{so}^2} - \frac{u^2}{2l_B^2}}$. The Laughlin-type wavefunction can be constructed as

$$\Psi(w_1, v_1; \dots; w_{N_x N_y}, v_{N_x N_y}) = (\det[w_i^{h_x} v_i^{h_z}])^q, \quad (13)$$

where q is an odd number; $\det[w_i^{h_x} v_i^{h_z}]$ represents the Slater determinant for the fully filled single particle states $w^{h_x} v^{h_z}$ and $0 \leq h_x \leq N_x - 1$ and $0 \leq h_z \leq N_z - 1$. The study of the topological properties of Eq. 13 will be deferred to a later publication.

In conclusion, we have generalized LLs into 3d and 4d in the Landau-like gauge by coupling spatially dependent SO coupling with harmonic potentials. This method can be generalized to arbitrary dimensions by replacing the Pauli-matrices with the Γ -matrices in corresponding dimensions. These high dimensional LLs exhibit spatial separation of helical or chiral fermion modes with opposite helicities, which give rises to gapless helical or chiral boundary modes. The 4d LLs give rise to the quantized non-linear electromagnetic responses as a spatially separated (3+1)d chiral anomaly.

CW thanks J. Hirsch for helpful discussions. YL thanks N. P. Ong and Z. Wang for helpful discussions and the Inamori Fellowship. YL and CW are supported by the NSF DMR-1105945 and AFOSR FA9550-11-1-0067(YIP) and SCZ is supported by the NSF under grant numbers DMR-0904264.

-
- [1] X.-L. Qi and S.-C. Zhang, Phys. Today **63**, 33 (2010).
 - [2] X.-L. Qi and S.-C. Zhang, Rev. Mod. Phys. **83**, 1057 (2011).
 - [3] M. Hasan and C. Kane, Rev. Mod. Phys. **82**, 3045 (2010).
 - [4] D.-J. Thouless, M. Kohmoto, M.-P. Nightingale, and M. den Nijs, Phys. Rev. Lett. **49**, 405 (1982).
 - [5] M. Kohmoto, Ann. Phys. **160**, 343 (1985).
 - [6] F. D. M. Haldane, Phys. Rev. Lett. **61**, 2015 (1988).
 - [7] S.-C. Zhang and J.-P. Hu, Science **294**, 823 (2001).
 - [8] X.-L. Qi, T.-L. Hughes, and S.-C. Zhang, Phys. Rev. B **78**, 195424 (2008).

- [9] C.-L. Kane and E.-J. Mele, Phys. Rev. Lett. **95**, 146802 (2005).
- [10] B.-A. Bernevig and S.-C. Zhang, Phys. Rev. Lett. **96**, 106802 (2006).
- [11] B.-A. Bernevig, T.-L. Hughes, and S.-C. Zhang, Science **314**, 1757 (2006).
- [12] L. Fu and C.-L. Kane, Phys. Rev. B **76**, 045302 (2007).
- [13] J.-E. Moore and L. Balents, Phys. Rev. B **75**, 121306 (2007).
- [14] R. Roy, New J. Phys. **12**, 065009 (2010).
- [15] M. König *et al.*, Science **318**, 766 (2007).
- [16] D. Hsieh *et al.*, Nature **452**, 970 (2008).
- [17] H. Zhang *et al.*, Nature Phys. **5**, 438 (2009).
- [18] Y. Xia *et al.*, Nature Phys. **5**, 398 (2009).
- [19] Y. Chen *et al.*, Science **325**, 178 (2009).
- [20] X.-L. Qi, T.-L. Hughes, S. Raghu, and S.-C. Zhang, Phys. Rev. Lett. **102**, 187001 (2009).
- [21] A. Kitaev, *American Institute of Physics Conference Series*, Vol. 1134, 22-30 (2009).
- [22] S. Ryu, A. Schnyder, A. Furusaki, and A. Ludwig, New J. Phys. **12**, 065010 (2010).
- [23] D. Karabali and V. Nair, Nucl. Phys. B **641**, 533 (2002).
- [24] H. Elvang and J. Polchinski, Comptes Rendus Physique **4**, 405 (2003).
- [25] B.-A. Bernevig, J. Hu, N. Toumbas, and S.-C. Zhang, Phys. Rev. Lett. **91**, 236803 (2003).
- [26] K. Hasebe, Symmetry, Integrability and Geometry: Methods and Applications **6**, (2010).
- [27] J. M. Edge, J. Tworzydło, and C. W. J. Beenakker, Arxiv preprint arXiv:1206.0099 (2012).
- [28] Y. Li and C. Wu, Arxiv preprint arXiv:1103.5422 (2011).
- [29] Y. Li, K. Intriligator, Y. Yu, and C. Wu, Phys. Rev. B **85**, 085132 (2012).
- [30] Y. Li, X. Zhou, and C. Wu, Phys. Rev. B **85**, 125122 (2012).
- [31] J. E. Hirsch, private communication and Arxiv preprint arXiv:1106.5311 (2011).
- [32] R. Jackiw and C. Rebbi, Phys. Rev. D **13**, 3398 (1976).
- [33] A. J. Niemi and G. W. Semenoff, Phys. Rep. **135**, 99 (1986).
- [34] E. H. Rezayi and F. D. M. Haldane, Phys. Rev. B **50**, 17199 (1994).